

## LECTURE 9

### 1. STANDARD FILTRATIONS

Last time we proved  $\mathcal{O}$  has enough projectives and injectives. Hence for  $M, N \in \mathcal{O}$ , the vector space  $\text{Ext}_{\mathcal{O}}^i(M, N)$ ,  $i \geq 0$  is well-defined. It can be calculated in either of the following methods:

- Choose a projective resolution  $P^\bullet \rightarrow M$  and consider the chain complex  $\text{Hom}_{\mathcal{O}}(P^\bullet, N)$ . Then  $\text{Ext}_{\mathcal{O}}^i(M, N)$  is the  $i$ -th cohomology of this complex.
- Choose an injective resolution  $N \rightarrow I^\bullet$  and consider the chain complex  $\text{Hom}_{\mathcal{O}}(M, I^\bullet)$ . Then  $\text{Ext}_{\mathcal{O}}^i(M, N)$  is the  $i$ -th cohomology of this complex.

We will prove the following result:

**Theorem-Definition 1.1.** *For an object  $M \in \mathcal{O}$ , the following conditions are equivalent:*

- (a) *The object  $M$  admits a finite filtration such that the subquotients are isomorphic to Verma modules.*
- (b) *For any weight  $\mu$  and  $i > 0$ ,  $\text{Ext}_{\mathcal{O}}^i(M, M_\mu^\vee) = 0$ .*
- (c) *For any weight  $\mu$ ,  $\text{Ext}_{\mathcal{O}}^1(M, M_\mu^\vee) = 0$ .*

A filtration as in (a) is called a **standard filtration** of  $M$ . Let  $\mathcal{O}^\Delta \subset \mathcal{O}$  be the full subcategory of objects admitting standard filtrations.

Via contragradient duality, the above result is equivalent to:

**Theorem-Definition 1.2.** *For an object  $M \in \mathcal{O}$ , the following conditions are equivalent:*

- (a) *The object  $M$  admits a finite filtration such that the subquotients are isomorphic to dual Verma modules.*
- (b) *For any weight  $\lambda$  and  $i > 0$ ,  $\text{Ext}_{\mathcal{O}}^i(M_\lambda, M) = 0$ .*
- (c) *For any weight  $\lambda$ ,  $\text{Ext}_{\mathcal{O}}^1(M_\lambda, M) = 0$ .*

A filtration as in (a) is called a **costandard filtration** of  $M$ . Let  $\mathcal{O}^\nabla \subset \mathcal{O}$  be the full subcategory of objects admitting costandard filtrations.

As a particular case, we obtain

**Corollary 1.3.** *For weights  $\lambda, \mu$  and  $i > 0$ , we have  $\text{Ext}_{\mathcal{O}}^i(M_\lambda, M_\mu^\vee) = 0$ .*

We will also prove the following result.

**Theorem 1.4.** *Every projective object of  $\mathcal{O}$  admits a standard filtration.*

To prove these results, we need some preparations.

**Lemma 1.5.** *The generalized Verma modules<sup>1</sup>  $M_{\lambda, n}$  admit standard filtrations.*

*Proof.* Recall  $M_{\lambda, n} \simeq \text{ind}_{\mathfrak{b}}^{\mathfrak{g}}(U(\mathfrak{b})/I'_{\lambda, n})$ , where  $I'_{\lambda, n}$  is the left ideal of  $U(\mathfrak{b})$  generated by the following elements:

- The element  $t - \lambda(t)$  for any  $t \in \mathfrak{t}$ ;

---

*Date:* Apr 22, 2024.

<sup>1</sup>See [Proof of Lemma 4.10, Lecture 8].

- The element  $x_1 x_2 \cdots x_n$  for  $x_i \in \mathfrak{n}^+$ .

By considering the weights, the finite-dimensional  $\mathfrak{b}$ -module  $U(\mathfrak{b})/I'_{\lambda,n}$  has a filtration such that the subquotients are 1-dimensional. Since the functor  $\text{ind}_{\mathfrak{b}}^{\mathfrak{g}}$  is exact, we obtain a standard filtration of  $M_{\lambda,n}$ . □

**Lemma 1.6.** *Let  $M \in \mathcal{O}^\Delta$  be an object admitting a standard filtration  $F^{\leq \bullet} M$  with length  $m$ . Let  $v \in M$  be a nonzero highest weight vector with weight  $\lambda$ , and  $M_\lambda \rightarrow M$  be the unique  $\mathfrak{g}$ -linear map sending  $v_\lambda$  to  $v$ . Let  $i$  be the smallest index such that the image  $\text{Im}(f)$  is contained in  $F^{\leq i} M$ . Then:*

- (1) *The composition  $M_\lambda \rightarrow M \rightarrow \text{gr}^i M$  is an isomorphism. In particular,  $M_\lambda \rightarrow M$  is injective.*
- (2) *The quotient  $M/M_\lambda$  admits a standard filtration with length  $m - 1$ .*

*Proof.* By assumption  $\text{gr}^i M$  is a Verma module  $M_\mu$  and the composition  $M_\lambda \rightarrow M \rightarrow \text{gr}^i M$  is nonzero. This implies  $\lambda \leq \mu$ . Since  $\lambda$  is a highest weight of  $M$ . We get  $\lambda = \mu$ . This implies  $M_\lambda \rightarrow M \rightarrow \text{gr}^i M$  is an isomorphism because it preserves the nonzero highest weight vectors. This proves (1).

For (2), we have a short exact sequence  $0 \rightarrow F^{\leq i-1} M \rightarrow M/M_\lambda \rightarrow M/F^{\leq i} M \rightarrow 0$ . By assumption,  $F^{\leq i-1} M$  has a standard filtration of length  $i - 1$ , and  $M/F^{\leq i} M$  has one of length  $m - i$ . It follows that  $M/M_\lambda$  admits a standard filtration with length  $(i - 1) + (m - i) = m - 1$ . □

**Lemma 1.7.** *For direct sum decomposition  $M = M_1 \oplus M_2$  in  $\mathcal{O}$ , if  $M$  admits a standard filtration, so do  $M_1$  and  $M_2$ .*

*Proof.* We use induction on the length of the standard filtration of  $M$ . When the length is 0, the claim is trivial. If the length is  $m > 0$ , then  $M \neq 0$ . Without loss of generality, we can assume  $M_1$  contains a nonzero highest weight vector  $v$  of  $M$  with weight  $\lambda$ . By Lemma 1.6, we have injections  $M_\lambda \hookrightarrow M_1 \hookrightarrow M$  such that  $M/M_\lambda$  admits a standard filtration of length  $m - 1$ . Since  $M/M_\lambda \simeq M_1/M_\lambda \oplus M_2$ , by induction hypothesis, we have  $M_1/M_\lambda, M_2 \in \mathcal{O}^\Delta$ . It follows that we also have  $M_1 \in \mathcal{O}^\Delta$ . □

*Proof of Theorem 1.4.* In the proof of [Theorem 4.3, Lecture 8], we proved any object in  $\mathcal{O}$  is a quotient of a direct sum of some  $M_{\lambda,n}$ . In general, if a projective object is a quotient of another object, then it is a direct summand of the latter. It follows that any projective object is a direct summand of a direct sum of some  $M_{\lambda,n}$ . Then the claim follows from Lemma 1.5 and Lemma 1.7. □[Theorem 1.4]

**Lemma 1.8.** *Let  $0 \rightarrow K \rightarrow M \rightarrow N \rightarrow 0$  be a short exact sequence such that  $N \in \mathcal{O}^\Delta$ . Then  $M \in \mathcal{O}^\Delta$  iff  $K \in \mathcal{O}^\Delta$ .*

**Warning 1.9.**  *$K, M \in \mathcal{O}^\Delta$  does not imply  $N \in \mathcal{O}^\Delta$ . This can be seen from the  $\mathfrak{sl}_2$ -case and the short exact sequence  $0 \rightarrow M_{-l-2} \rightarrow M_l \rightarrow L_l \rightarrow 0$ .*

*Proof.* The “if” part is obvious. For the “only if” part, let  $M \in \mathcal{O}^\Delta$ . Using induction, it is easy to reduce to the case when  $N$  is a Verma module  $M_\mu$ . Let  $M_\lambda \rightarrow M$  be as in Lemma 1.6. Then  $M/M_\lambda \in \mathcal{O}^\Delta$ . Note that the composition  $M_\lambda \rightarrow M \rightarrow M_\mu$  is either 0 or an isomorphism by considering the weights. If this is the zero map, then  $M/M_\lambda \rightarrow M_\mu$  is still a surjection and we can finish the proof by using induction. Otherwise  $M \simeq K \oplus M_\mu$  and the claim follows from Lemma 1.7.

□

*Proof of Theorem-Definition 1.1.* We first prove (a)  $\Rightarrow$  (b). We prove by induction on  $i$ . For any fixed  $i$ , using the long exact sequences

$$\cdots \rightarrow \text{Ext}_{\mathcal{O}}^i(\mathbb{F}^{\leq k-1}M, M_{\mu}^{\vee}) \rightarrow \text{Ext}_{\mathcal{O}}^i(\mathbb{F}^{\leq k}M, M_{\mu}^{\vee}) \rightarrow \text{Ext}_{\mathcal{O}}^i(\text{gr}^k M, M_{\mu}^{\vee}) \rightarrow \cdots,$$

we only need to show  $\text{Ext}_{\mathcal{O}}^i(\text{gr}^k M, M_{\mu}^{\vee}) = 0$  for any  $k$ . By assumption,  $\text{gr}^k M \simeq M_{\lambda}$  for some weight  $\lambda$ . The case  $i = 1$  is just [Lemma 3.16, Lecture 8]. For  $i > 1$ , let  $0 \rightarrow N \rightarrow P \rightarrow M_{\lambda} \rightarrow 0$  be a short exact sequence such that  $P$  is projective. We have a long exact sequence

$$\cdots \rightarrow \text{Ext}_{\mathcal{O}}^{i-1}(P, M_{\mu}^{\vee}) \rightarrow \text{Ext}_{\mathcal{O}}^{i-1}(N, M_{\mu}^{\vee}) \rightarrow \text{Ext}_{\mathcal{O}}^i(M_{\lambda}, M_{\mu}^{\vee}) \rightarrow \text{Ext}_{\mathcal{O}}^i(P, M_{\mu}^{\vee}) \rightarrow \cdots.$$

Since  $P$  is projective, we have  $\text{Ext}_{\mathcal{O}}^{i-1}(N, M_{\mu}^{\vee}) \simeq \text{Ext}_{\mathcal{O}}^i(M_{\lambda}, M_{\mu}^{\vee})$ . By Theorem 1.4 and Lemma 1.8,  $N \in \mathcal{O}^{\Delta}$ . Hence by induction hypothesis  $\text{Ext}_{\mathcal{O}}^{i-1}(N, M_{\mu}^{\vee}) \simeq 0$  as desired.

(b)  $\Rightarrow$  (c) is obvious.

It remains to show (c)  $\Rightarrow$  (a).

We prove by induction on  $\text{length}(M)$ . The case  $\text{length}(M) = 0$  is obvious. If  $\text{length}(M) > 0$ , let  $\lambda$  be a highest weight vector of  $M$  and  $n := \dim(M^{\text{wt}=\lambda})$ . Then there is a nonzero  $\mathfrak{g}$ -linear map  $M_{\lambda}^{\oplus n} \rightarrow M$ . Let<sup>2</sup>  $N_1$  and  $N_2$  be the kernel and cokernel of this map, and  $M' \neq 0$  be the image of it. We have short exact sequences

$$\begin{aligned} 0 \rightarrow N_1 \rightarrow M_{\lambda}^{\oplus n} \rightarrow M' \rightarrow 0, \\ 0 \rightarrow M' \rightarrow M \rightarrow N_2 \rightarrow 0. \end{aligned}$$

They induce long exact sequences

$$\begin{aligned} 0 \rightarrow \text{Hom}_{\mathcal{O}}(M', M_{\mu}^{\vee}) \rightarrow \text{Hom}_{\mathcal{O}}(M_{\lambda}^{\oplus n}, M_{\mu}^{\vee}) \rightarrow \text{Hom}_{\mathcal{O}}(N_1, M_{\mu}^{\vee}) \rightarrow \text{Ext}_{\mathcal{O}}^1(M', M_{\mu}^{\vee}) \rightarrow \cdots \\ \cdots \rightarrow \text{Hom}_{\mathcal{O}}(M, M_{\mu}^{\vee}) \rightarrow \text{Hom}_{\mathcal{O}}(M', M_{\mu}^{\vee}) \rightarrow \text{Ext}_{\mathcal{O}}^1(N_2, M_{\mu}^{\vee}) \rightarrow \text{Ext}_{\mathcal{O}}^1(M, M_{\mu}^{\vee}) \rightarrow \\ \rightarrow \text{Ext}_{\mathcal{O}}^1(M', M_{\mu}^{\vee}) \rightarrow \text{Ext}_{\mathcal{O}}^2(N_2, M_{\mu}^{\vee}) \rightarrow \cdots \end{aligned}$$

By assumption,  $\text{Ext}_{\mathcal{O}}^1(M, M_{\mu}^{\vee}) = 0$ . We claim  $\text{Ext}_{\mathcal{O}}^1(N_2, M_{\mu}^{\vee}) = 0$ . Indeed:

- If  $\lambda \neq \mu$ , then  $\text{Hom}_{\mathcal{O}}(M_{\lambda}^{\oplus n}, M_{\mu}^{\vee}) = 0$  ([Lemma 3.14, Lecture 8]). By the first sequence,  $\text{Hom}_{\mathcal{O}}(M', M_{\mu}^{\vee}) = 0$ . By the second sequence,  $\text{Ext}_{\mathcal{O}}^1(N_2, M_{\mu}^{\vee}) = 0$ .
- If  $\lambda = \mu$ , note that  $\mu$  is a highest weight of both  $M'$  and  $M$  and by construction,  $(M')^{\text{wt}=\mu} \simeq M^{\text{wt}=\mu}$ . Hence we have

$$\text{Hom}_{\mathcal{O}}(M', M_{\mu}^{\vee}) \simeq ((M')^{\text{wt}=\mu})^* \simeq (M^{\text{wt}=\mu})^* \simeq \text{Hom}_{\mathcal{O}}(M, M_{\mu}^{\vee}).$$

By the second sequence,  $\text{Ext}_{\mathcal{O}}^1(N_2, M_{\mu}^{\vee}) = 0$ .

Note that  $\text{length}(N_2) < \text{length}(M)$ . By the induction hypothesis,  $N_2 \in \mathcal{O}^{\Delta}$ . Using (a)  $\Rightarrow$  (b), we get  $\text{Ext}_{\mathcal{O}}^2(N_2, M_{\mu}^{\vee}) = 0$ . By the second sequence,  $\text{Ext}_{\mathcal{O}}^1(M', M_{\mu}^{\vee}) = 0$ . Now we have two cases:

- If  $N_2 \neq 0$ , then  $\text{length}(M') < \text{length}(M)$ . By the induction hypothesis,  $M' \in \mathcal{O}^{\Delta}$ . Then  $M \in \mathcal{O}^{\text{length}}$  because it is an extension of  $N_2$  by  $M'$ .
- If  $N_2 = 0$ , then  $M \simeq M'$ . An argument similar to that in the last paragraph implies  $\text{Hom}_{\mathcal{O}}(N_1, M_{\mu}^{\vee}) = 0$ . Note that  $\mu$  can be any weight. This forces  $N_1 = 0$  and therefore  $M \simeq M_{\lambda}^{\oplus n}$ . Then it is clear  $M \in \mathcal{O}^{\Delta}$ .

□[Theorem-Definition 1.1]

**Lemma 1.10.** *Let  $M \in \mathcal{O}_{\chi}$  and  $M' \in \mathcal{O}_{\chi'}$  such that  $\chi \neq \chi'$ . Then  $\text{Ext}^i(M, N) = 0$  for any  $i \geq 0$ .*

<sup>2</sup>The following language can be rewritten in the language of spectral sequences.

*Proof.* Let  $P^\bullet \rightarrow M$  be a projective resolution of  $M$ . We can replace each  $P^{-n}$  by its image under the functor  $\mathcal{O} \rightarrow \mathcal{O}_\chi$  and obtain a projective resolution of  $M$  contained in  $\mathcal{O}_\chi$ . Now the claim follows from the case  $i = 0$ .  $\square$

*Exercise 1.11.* This is **Homework 4, Problem 4**. Let  $\lambda$  be a weight. Prove:

- (1) If  $M \in \mathcal{O}$  such that  $\text{wt}(M) \cap \{\mu \mid \mu \geq \lambda\} = \emptyset$ , then  $\text{Ext}_{\mathcal{O}}^i(M, M_\lambda^\vee) = 0$  for  $i \geq 0$ <sup>3</sup>.
- (2)  $\text{Ext}_{\mathcal{O}}^i(L_\lambda, M_\lambda^\vee) = 0$  for  $i > 0$ .
- (3) Combining (1) and (2), deduce  $\text{Ext}_{\mathcal{O}}^i(M_\lambda, L_\mu) = 0$  and  $\text{Ext}_{\mathcal{O}}^i(L_\mu, M_\lambda^\vee) = 0$  for  $i > 0$  and  $\mu \not\geq \lambda$ .

## 2. BGG RECIPROCITY

The following result follows from Theorem-Definition 1.1 by dévissage.

**Proposition-Definition 2.1.** *Let  $M \in \mathcal{O}^\Delta$ , then for any standard filtration of  $M$ , the multiplicity of  $M_\lambda$  in the subquotients does not depend on the filtration. We denote this number by  $(M : M_\lambda)$ . Moreover, we have*

$$(M : M_\lambda) \simeq \dim(\text{Hom}_{\mathcal{O}}(M, M_\lambda^\vee)).$$

**Theorem 2.2** (BGG reciprocity). *For weights  $\lambda, \mu \in \mathfrak{t}^*$ , we have*

$$(P_\mu : M_\lambda) = [M_\lambda^\vee : L_\mu] = [M_\lambda : L_\mu].$$

*Proof.* The last identity follows from  $L_\mu^\vee \simeq L_\mu$ . The first one follows from Proposition-Definition 2.1 and [Corollary 4.9, Lecture 8].  $\square$

*Remark 2.3.* The previous discussions on standard filtrations and BGG reciprocity can be axiomized using the language of **highest weight categories**. See [CPS].

*Remark 2.4.* In  $\mathcal{O}$ , or any highest weight category, if an object admits both a standard filtration and a costandard filtration, then it is called a **tilting object**. One can show the set of indecomposable tilting objects is bijective to the set of irreducible objects. Tilting objects play important roles in representation theory. For a geometry-oriented<sup>4</sup> introduction, see [BBM].

## 3. TRANSLATION FUNCTORS

**Construction 3.1.** *Let  $V$  be a finite-dimensional  $\mathfrak{g}$ -module. Consider the functor*

$$\mathfrak{g}\text{-mod} \rightarrow \mathfrak{g}\text{-mod}, M \mapsto V \otimes M,$$

*where (recall) the  $\mathfrak{g}$ -module structure on  $V \otimes M$  is given by the Leibniz rule. It is easy to see this functor preserves  $\mathcal{O}$ . We denote the obtained functor by  $T_V : \mathcal{O} \rightarrow \mathcal{O}$ . Note that  $T_V$  is exact.*

**Lemma 3.2.** *The functor  $T_V : \mathcal{O} \rightarrow \mathcal{O}$  commutes with contragredient duality.*

*Proof.* For any finite dimensional  $\mathfrak{g}$ -module, we have  $V^\vee \simeq V$  because  $L_\lambda^\vee \simeq L_\lambda$ .  $\square$

**Lemma 3.3.** *The functor  $T_{V^*}$  is both left and right adjoint to  $T_V$ . In particular,  $T_V$  preserves both projectives and injectives.*

<sup>3</sup>Hint: Step 1: reduce to the case  $M = L_\mu$  with  $\varpi(\mu) = \varpi(\lambda)$  and  $\mu \not\geq \lambda$ . Step 2: consider  $0 \rightarrow K \rightarrow M_\mu \rightarrow L_\mu \rightarrow 0$  and note that  $\text{wt}(K) < \mu$ .

<sup>4</sup>You should ask experts in representation theory for other good references.

*Proof.* We have the unit and pairing maps  $k \rightarrow V \otimes V^*$  and  $V^* \otimes V \rightarrow k$ , which induce natural transformations  $T_k \rightarrow T_{V \otimes V^*}$  and  $T_{V^* \otimes V} \rightarrow T_k$ . Note that  $T_k \simeq \text{Id}$  and  $T_{V \otimes V^*} \simeq T_V \circ T_{V^*}$ . Hence we obtain natural transformations  $\text{Id} \rightarrow T_V \circ T_{V^*}$  and  $T_{V^*} \circ T_V \rightarrow \text{Id}$ . Now the duality data between  $V$  and  $V^*$  are translated exactly to the adjunction data between  $T_V$  and  $T_{V^*}$ .  $\square$

*Remark 3.4.* It follows formally that  $T_{V^*}$  and  $T_V$  are adjoint in the derived sense, i.e.  $\text{Ext}^i(T_V(M), N) \simeq \text{Ext}^i(M, T_{V^*}(N))$ .

**Lemma 3.5.** *For any weight  $\lambda$ , the module  $T_V(M_\lambda)$  admits a standard filtration  $F^{\leq k}(T_V(M_\lambda))$  such that the highest weights of  $\text{gr}^k(T_V(M_\lambda))$  is (non-strictly) decreasing in  $k$ . Moreover,*

$$(T_V(M_\lambda) : M_\mu) = \dim V^{\text{wt}=\mu-\lambda}.$$

*Proof.* Follows from the *projection formula*

$$\text{ind}_{\mathfrak{b}}^{\mathfrak{g}}(V_1) \otimes V_2 \simeq \text{ind}_{\mathfrak{b}}^{\mathfrak{g}}(V_1 \otimes V_2), \quad V_1 \in \mathfrak{b}\text{-mod}, V_2 \in \mathfrak{g}\text{-mod}.$$

Indeed, any finite-dimensional weight  $\mathfrak{b}$ -module, such as  $V \otimes k_\lambda$ , admits a finite filtration whose subquotients are 1-dimensional modules with decreasing weights.  $\square$

**Construction 3.6.** *Let  $\chi_1, \chi_2$  be two central characters. For any finite-dimensional  $\mathfrak{g}$ -module  $V$ , consider the composition*

$$T_{\chi_1, V, \chi_2} : \mathcal{O}_{\chi_1} \rightarrow \mathcal{O} \xrightarrow{T_V} \mathcal{O} \rightarrow \mathcal{O}_{\chi_2}.$$

*We call such functors the **translation functors**. It follows that  $T_{\chi_1, V, \chi_2}$  is exact, and is both left and right adjoint to  $T_{\chi_2, V^*, \chi_1}$ .*

**Construction 3.7.** *Let  $\lambda$  and  $\mu$  be weights such that  $\mu - \lambda$  is integral. Then the  $W$ -orbit (for the linear action)  $W(\mu - \lambda)$  contains a unique dominant integral weight  $\nu$ . Consider the finite-dimensional  $\mathfrak{g}$ -module  $L_\nu$ . We write*

$$T_\lambda^\mu := T_{\varpi(\lambda), L_\nu, \varpi(\mu)} : \mathcal{O}_{\varpi(\lambda)} \rightarrow \mathcal{O}_{\varpi(\mu)}.$$

*Note that  $w_0(-\nu)$  is also dominant and integral. By definition, we have*

$$T_\mu^\lambda = T_{\varpi(\mu), L_{-w_0(\nu)}, \varpi(\lambda)},$$

*which is both left and right adjoint to  $T_\lambda^\mu$  because  $L_{-w_0(\nu)} \simeq L_\nu^*$ <sup>5</sup>*

Recall the following definition:

**Definition 3.8.** Let  $(E, \Phi)$  be the root system of  $\mathfrak{g}$ . For  $\lambda, \mu \in E$ , we say they belong to the same **dot-Weyl facet** if the signs of  $\langle \lambda + \rho, \check{\alpha} \rangle$  and  $\langle \mu + \rho, \check{\alpha} \rangle$  are the same for any  $\alpha \in \Phi$ .

For  $\lambda \in E$ , let  $F_\lambda$  be the dot-Weyl facet containing it.

**Definition 3.9.** For a dot-Weyl facet  $F_\lambda$ , its **upper closure**  $F_\lambda^+$  is the subset of  $\mu \in E$  such that

$$\langle \mu + \rho, \check{\alpha} \rangle \begin{cases} > 0 & \text{if } \langle \lambda + \rho, \check{\alpha} \rangle > 0, \\ = 0 & \text{if } \langle \lambda + \rho, \check{\alpha} \rangle = 0, \\ \leq 0 & \text{if } \langle \lambda + \rho, \check{\alpha} \rangle < 0 \end{cases}$$

<sup>5</sup>Here  $L_\nu^*$  is the usual linear dual of  $L_\nu$ . It is not the contragredient dual. Note that  $w_0(\nu)$  is indeed the lowest weight of  $L_\nu$ .

*Remark 3.10.* When equipped with the standard topology,  $F_\lambda$  is a locally closed subset of  $E$ , and both the closure  $\overline{F_\lambda}$  and the upper closure  $F_\lambda^+$  are unions of dot-Weyl facets.

Note that each dot-Weyl facet is contained in the upper closure of a unique **dot-Weyl chamber**, i.e., open dot-Weyl facet. Also,  $\lambda$  is dot-regular iff  $F_\lambda$  is a dot-Weyl chamber.

**Example 3.11.** For  $\mathfrak{sl}_2$  and the coordinate  $l := \langle \lambda, \check{\alpha} \rangle$ , there are three facets:  $(-\infty, -1)$ ,  $\{-1\}$  and  $(-1, \infty)$ . Note that  $\{-1\}$  is contained in the upper closure of  $(-\infty, -1)$ .

We have the following theorem. For complete proofs and its generalization to non-integral weights, see [H, Sect. 7]<sup>6</sup>.

**Theorem 3.12.** *Let  $\lambda$  and  $\mu$  be dot-antidominant integral weights such that  $F_\mu \subset \overline{F_\lambda}$ . Then for any  $w \in W$*

$$T_\lambda^\mu(M_{w \cdot \lambda}) \simeq M_{w \cdot \mu}, \quad T_\lambda^\mu(M_{w \cdot \lambda}^\vee) \simeq M_{w \cdot \mu}^\vee$$

and

$$T_\lambda^\mu(L_{w \cdot \lambda}) \simeq \begin{cases} L_{w \cdot \mu} & \text{if } F_{w \cdot \mu} \subset F_{w \cdot \lambda}^+, \\ 0 & \text{otherwise.} \end{cases}$$

*Sketch.* Let  $\nu \in W(\mu - \lambda)$  be the unique dominant integral weight in this orbit. Write  $V := L_\nu$ . By Lemma 3.5,  $T_\lambda^\mu(M_{w \cdot \lambda})$  admits a standard filtration and the multiplicity

$$(T_\lambda^\mu(M_{w \cdot \lambda}) : M_{w' \cdot \mu}) = \dim V^{\text{wt}=w' \cdot \mu - w \cdot \lambda}.$$

The RHS is nonzero unless  $w' \cdot \mu - w \cdot \lambda \leq \nu$ . Now a combinatorial argument (see [H, Lemma 7.5]) shows the latter can happen only if  $w' \cdot \mu = w \cdot \mu$ , and in this case the multiplicity is 1 because  $w \cdot \mu - w \cdot \lambda = w(\mu - \lambda) \in W(\nu)$ . This implies  $T_\lambda^\mu(M_{w \cdot \lambda}) \simeq M_{w \cdot \mu}$ .

The statement for dual Verma modules follows from the contragredient duality.

Now consider the chain  $M_{w \cdot \lambda} \twoheadrightarrow L_{w \cdot \lambda} \hookrightarrow M_{w \cdot \lambda}^\vee$ . Since  $T_\lambda^\mu$  is exact, we obtain a chain  $M_{w \cdot \mu} \twoheadrightarrow T_\lambda^\mu(L_{w \cdot \lambda}) \hookrightarrow M_{w \cdot \mu}^\vee$ . This forces  $T_\lambda^\mu(L_{w \cdot \lambda}) \simeq L_{w \cdot \mu}$  or 0 ([Lemma 3.14, Lecture 8]).

It remains to pinpoint these two cases. Since the exact functor  $T_\lambda^\mu$  sends  $M_{w \cdot \lambda}$  to  $M_{w \cdot \mu}$ , there exists a unique composition factor  $L_{w' \cdot \lambda}$  of  $M_{w \cdot \lambda}$  such that  $T_\lambda^\mu(L_{w' \cdot \lambda}) \simeq L_{w \cdot \mu}$ . By the last paragraph, we must have  $w' \cdot \mu = w \cdot \mu$ . Now we have two cases:

- If  $F_{w \cdot \mu}$  is contained in the upper closure  $F_{w \cdot \lambda}^+$ , then a combinatorial argument shows  $w' \cdot \lambda = w \cdot \lambda$  and therefore  $T_\lambda^\mu(L_{w \cdot \lambda}) \simeq L_{w \cdot \mu}$  as desired.
- If  $F_{w \cdot \mu}$  is not contained in the upper closure  $F_{w \cdot \lambda}^+$ , then there exists a reflection  $s_\alpha$  such that  $s_\alpha w \cdot \lambda < w \cdot \lambda$  while  $s_\alpha w \cdot \mu = w \cdot \mu$ . By Verma's theorem, there exists a proper embedding  $M_{s_\alpha w \cdot \lambda} \subset M_{w \cdot \lambda}$  which is sent to an isomorphism by  $T_\lambda^\mu$ . Hence  $L_{w \cdot \lambda}$ , which is a quotient of  $M_{w \cdot \lambda} / M_{s_\alpha w \cdot \lambda}$ , is sent to 0 as desired.

□

The following result is a formal consequence of the above theorem:

**Theorem 3.13.** *Let  $\lambda$  and  $\mu$  be dot-antidominant integral weights such that  $F_\lambda = F_\mu$ . Then  $T_\lambda^\mu : \mathcal{O}_{\varpi(\lambda)} \rightarrow \mathcal{O}_{\varpi(\mu)}$  is an equivalence.*

*Proof.* By the previous theorem, the exact functor  $F := T_\lambda^\mu$  is left adjoint to the exact functor  $G := T_\mu^\lambda$ , and they induce a bijection between the sets of irreducible objects. In general, such adjoint functors are inverse to each other.

*Proof:* we only to show the adjunctions  $\text{Id} \rightarrow G \circ F$  and  $F \circ G \rightarrow \text{Id}$  are equivalences. We will prove the first equivalence and the second follows similarly. Note that for any object  $M$ ,  $M$  and  $G \circ F(M)$  have the same composition factors and multiplicities. Hence we only need to show  $M \rightarrow G \circ F(M)$  is injective. Since  $F$  sends nonzero objects to nonzero objects, we only need to

<sup>6</sup>See [G, Sect. 4.23] for a simplified proof in a special case.

show  $F(M) \rightarrow F \circ G \circ F(M)$  is injective. By the axiom of adjoint functors, this morphism has a left inverse, hence is indeed injective.  $\square$

*Remark 3.14.* The above theorems essentially reduce the study about any integral block  $\mathcal{O}_\chi$  to the principle block  $\mathcal{O}_{\varpi(0)}$ .

**Example 3.15.** Any dot-regular integral block  $\mathcal{O}_\chi$  is equivalent to the principle block  $\mathcal{O}_{\varpi(0)}$ . Indeed, this is the special case of the above theorem when  $\lambda = 0$  and  $\mu \in \varpi^{-1}(\chi)$  is dot-antidominant. Note that the equivalence  $T_0^\mu$  preserves (dual) Verma modules and irreducible modules.

**Example 3.16.** For  $\mathfrak{g} = \mathfrak{sl}_2$  and the coordinate  $l := \langle \lambda, \check{\alpha} \rangle$ , then  $\mathcal{O}_{\varpi(0)} \simeq \mathcal{O}_{\varpi(l)}$  for  $l \in \mathbb{Z}^{\geq 0}$  such that the short exact sequence  $0 \rightarrow M_{-2} \rightarrow M_0 \rightarrow L_0 \rightarrow 0$  is sent to  $0 \rightarrow M_{-l-2} \rightarrow M_l \rightarrow L_l \rightarrow 0$ .

On the other hand, the translation functor  $\mathcal{O}_{\varpi(0)} \rightarrow \mathcal{O}_{\varpi(-1)}$  sends this sequence to  $0 \rightarrow M_{-1} \rightarrow M_{-1} \rightarrow 0 \rightarrow 0$ . Note that  $L_{-2}$  is sent to  $L_{-1}$  and  $-1 \in F_{-2}^+$ , while  $L_0$  is sent to 0 and  $-1 \notin F_0^+$ .

**Construction 3.17.** Let  $\mu$  be any dot-antidominant integral weight. The functor  $T_{-\rho}^\mu : \mathcal{O}_{\varpi(-\rho)} \rightarrow \mathcal{O}_{\varpi(\mu)}$  is not covered by the above theorems (although its adjoint  $T_\mu^{-\rho}$  is). Recall the most singular block  $\mathcal{O}_{\varpi(-\rho)}$  is semi-simple and contains a unique irreducible object  $L_{-\rho}$ , which is both projective and injective ([Example 4.16, Lecture 7]). It follows that

$$\Xi_\mu := T_{-\rho}^\mu(L_{-\rho})$$

is both projective and injective.

*Exercise 3.18.* This is **Homework 4, Problem 5**. Let  $\mu$  be any dot-antidominant integral weight. Prove<sup>7</sup>:

- (1) For any  $w \in W$ ,  $(\Xi_\mu : M_{w\cdot\mu}) = 1$  and there is a surjection  $\Xi_\mu \twoheadrightarrow M_\mu$ .
- (2) For any  $w \in W$ ,  $(P_\mu : M_{w\cdot\mu}) \geq 1$  and there is a surjection  $P_\mu \twoheadrightarrow M_\mu$ .
- (3) There exists an isomorphism  $\Xi_\mu \simeq P_\mu$  compatible with the surjections to  $M_\mu$ .
- (4) For any  $w \in W$ ,  $[M_{w\cdot\mu} : L_\mu] = 1$ <sup>8</sup>.

*Remark 3.19.* For dot-antidominant weight  $\mu$ , the projective  $P_\mu$  is called the **big projective module**, which plays an important role in representation theory.

#### REFERENCES

- [BBM] Beilinson, A., R. Bezrukavnikov, and I. Mirkovic. Tilting Exercises. *Moscow Mathematical Journal* 4, no. 3 (2004): 547-557.
- [CPS] Scott, L., Brian Parshall, and E. Cline. Finite dimensional algebras and highest weight categories. (1988): 85-99.
- [G] Gaitsgory, Dennis. Course Notes for Geometric Representation Theory, 2005, available at <https://people.mpim-bonn.mpg.de/gaitsgde/267y/cat0.pdf>.
- [H] Humphreys, James E. Representations of Semisimple Lie Algebras in the BGG Category  $\mathcal{O}$ . Vol. 94. American Mathematical Soc., 2008.

<sup>7</sup>Hint: Lemma 3.5 for (1); Theorem 2.2 and [Corollary 4.15, Lecture 7] for (2). For (3), first find a surjection  $\Xi_\mu \twoheadrightarrow P_\mu$  then use (1) and (2).

<sup>8</sup>See [G, Proposition 4.20] for a different proof of this fact.